

Closed Expressions for Finite Transformations

Daniel Lee Wenger

We find a solution to the following simple problem. Given an $n \times n$ matrix θ

$$\theta = \begin{pmatrix} \theta_{11} & \theta_{12} & \theta_{13} & \cdot & \theta_{1n} \\ \theta_{21} & \theta_{22} & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot \\ \theta_{n1} & \cdot & \cdot & \cdot & \theta_{nn} \end{pmatrix} \quad (1)$$

where the θ_{ij} are complex numbers and where θ is diagonalizable, then find a closed expression for the exponentiated form $\mu = e^{i\theta}$ where

$$\mu = e^{i\theta} = 1 + i\theta + \frac{(i\theta)^2}{2!} + \frac{(i\theta)^3}{3!} + \dots \quad (2)$$

By the Cayley-Hamilton theorem, the $n \times n$ matrix θ satisfies an n^{th} degree polynomial equation. Consequently, μ may be expressed in terms of a power series of degree $n-1$ in θ

$$\mu = \sum_{i=0}^{n-1} a_i \theta^i \quad (3)$$

The a_i are functions of the invariants of θ and the problem is to find these functions.

Now define the quantity

$$T_i \equiv \text{tr}(\mu \theta^i) \quad (4)$$

The trace is invariant under the transformation that diagonalizes θ and μ . Let the diagonal form of θ be $\bar{\theta}$ and the diagonal form of μ be $\bar{\mu}$, then

$$T_i = \text{tr}(\bar{\mu}\bar{\theta}^i) \quad (5)$$

Also, using (3) and (4) we have

$$T_i = \sum_{j=0}^{n-1} a_j \text{tr}(\theta^{i+j}) = \sum_{j=0}^{n-1} a_j \text{tr}(\bar{\theta}^{i+j}) \quad (6)$$

Let

$$A_{ij} \equiv \text{tr}(\bar{\theta}^{i+j}) \quad i, j = 0, 1, 2 \dots n-1 \quad (7)$$

Then

$$T_i = \sum_{j=0}^{n-1} A_{ij} a_j \quad i = 0, 1, 2 \dots n-1 \quad (8)$$

is a linear system of equations for a_j . Assuming that the determinant $|A| \neq 0$, the inverse to A exists and we have

$$a_i = \sum_{j=0}^{n-1} A_{ij}^{-1} T_j \quad (9)$$

As an example we find the 2 x 2 representation of SU_2 . θ is hermitian and traceless. Consequently, $\bar{\theta}$ has the form

$$\bar{\theta} = \begin{pmatrix} \phi/2 & 0 \\ 0 & -\phi/2 \end{pmatrix} \quad (10)$$

where ϕ is real, $\bar{\mu}$ has the form

$$\bar{\mu} = \begin{pmatrix} e^{i\phi/2} & 0 \\ 0 & e^{-i\phi/2} \end{pmatrix} \quad (11)$$

Computing T_i and A_{ij} we get

$$T_0 = 2 \cos \frac{\phi}{2} \quad T_1 = \frac{\phi}{2} \sin \frac{\phi}{2} \quad (12)$$

$$A = \begin{pmatrix} 2 & 0 \\ 0 & \phi^2/2 \end{pmatrix} \quad (13)$$

Inverting A we get

$$A^{-1} = \begin{pmatrix} 1/2 & 0 \\ 0 & 2/\phi^2 \end{pmatrix} \quad (14)$$

and solving for a_i from (9)

$$a_0 = \cos \frac{\phi}{2} \quad a_1 = \frac{i}{\phi/2} \sin \frac{\phi}{2} \quad (15)$$

we arrive at the familiar form

$$e^{i\theta} = \cos \frac{\phi}{2} + \frac{i}{\phi/2} \sin \frac{\phi}{2} \theta \quad (16)$$

where ϕ is the angle of rotation.